

1st Trimester OB US Technologist Worksheet

St Vincents Riverside Southside Clay St Johns Imaging Center Arlington ER Westside ER Beach ER
Optimal Forbes Southside Clay Mandarin Westside St Johns Town Center Orange Park

Patient Name: _____ MMI: _____ Age: _____

History/Symptoms: _____

Pelvic Pain	Nausea	Vaginal Bleeding	Vaginal Discharge	Trauma	Confirm Pregnancy
R L	Vomiting	Heavy Bleeding	Leaking Fluid	Prior Ectopic	Assess Fetal Viability
LMP	G	# Spont Abort:	Preg	Positive	Pending
	P	# Elect Abort:	Test	Not Ordered	Quant
					TA Scan
					EV Scan

Gestational Sac	Y	N	MSD	mm	EGA	EGA	Estimated Date
Yolk Sac	Y	N	CRL	mm	(by LMP)	(by AUA)	of Delivery
Fetal Pole	Y	N	Hrt Rate	bpm	wk d	wk d	/ /
Cardiac Activity	Y	N					

UTERUS	x	x	cm	anteverted	anteflexed	Cervix
				retroverted	retroflexed	closed opened
RT OVARY/ADNEXA	x	x	cm	flow present	flow absent	<i>poorly visualized</i>
LT OVARY/ADNEXA	x	x	cm	flow present	flow absent	<i>poorly visualized</i>

Other Findings: _____

FREE FLUID	none	trace	small	moderate	large
-------------------	------	-------	-------	----------	-------

Gestational sac should be seen by bHCG 3000 and EGA 4.5-5.0 wks.

Findings diagnostic of failed pregnancy: no heartbeat ≥ 7 mm CRL, no embryo ≥ 25 mm MSD, absent embryo with heartbeat ≥ 2 wks after US showed gestational sac without yolk sac, absence of embryo with heartbeat ≥ 11 days after US showed gestational sac with yolk sac.

Findings suspicious for failed pregnancy: no heartbeat 5 or 6 mm CRL, no embryo 16-24 mm MSD, absence of embryo with heartbeat 7–13 days after US showed gestational sac without yolk sac, absence of embryo with heartbeat 7–10 days after US showed gestational sac with yolk sac, absence of embryo 6 or more weeks after last menstrual period, empty amnion, > 7 mm yolk sac, < 5 mm difference between MSD and CRL.

Sonographer's Impression: _____

Sonographer's Name, Date & Time: _____ # Images _____

SN TV Probe: _____ HLD Completion Date/Time: _____ Performed By: _____